

Table 4

Modern pollen samples in each vegetation zone

Vegetation Zone ¹	Number of samples		
	Lake sediments	Moss polsters ²	Soils ³
<i>Picea sitchensis</i>	2	4	0
<i>Tsuga heterophylla</i>	5	32	0
<i>Abies amabilis</i>	10	3	0
<i>Tsuga mertensiana</i>	23	17	0
<i>Treeline/Alpine</i>	5	16	0
<i>Interior Valley</i>	1	7	0
<i>Mixed Evergreen</i>	2	0	0
<i>Mixed Conifer</i>	10	0	0
<i>Abies magnifica</i>	3	0	0
<i>Shrub Steppe</i>	5	4	78
<i>Juniperus</i>	4	4	0
<i>occidentalis</i>			
<i>Pinus ponderosa</i>	14	9	10
<i>Abies grandis</i>	7	14	8
<i>Abies lasiocarpa</i>	4	3	8

¹ Based on Franklin and Dyrness 1988² Heusser 1969, 1973, 1978a, 1978b³ Mack and Bryant 1974, Mack et al. 1978

Minckley, T. and Whitlock, C. (2000) Spatial variation of modern pollen in Oregon and southern Washington, USA. *Review of Palaeobotany and Palynology* 112: 97-123.